

### **What is Nanotechnology?**

Nanotechnology involves the manipulation of materials and the creation of structures and systems that exist at the scale of atoms and molecules. One nanometer (nm) is one thousandth of a micrometer ( $\mu\text{m}$ ), one millionth of a millimeter (mm) and one billionth of a meter (m). If you were to imagine that a nanometer is represented by a person, a red blood cell would be more than 4 miles long! The properties of nanoscale materials (measuring  $<100\text{nm}$ ) differ significantly from larger scales. Nanomaterials introduce new and often heightened risks of toxicity that remain poorly understood.

### **Nanoparticles are already being used (unlabelled) in hundreds of consumer products including sunscreens and cosmetics**

Our research demonstrates that nanoparticles have entered just about every personal care product on the market, including deodorant, soap, toothpaste, shampoo, hair conditioner, sunscreen, anti-wrinkle cream, moisturizer, foundation, face powder, lipstick, blush, eye shadow, nail polish, perfume and after-shave lotion. Nano-scale titanium dioxide and zinc is widely used in sunscreens. While we don't know how widespread the use of nanoparticles is in sunscreens on the U.S. market – in Australia estimates in 2006 showed that 70% of titanium dioxide sunscreens and 30% of zinc sunscreens sold in Australia contain 'manufactured' nanoparticles.<sup>i</sup> Major cosmetics manufacturers including Revlon, L'Oreal, Lancôme, Avon and The Body Shop also use nanoparticles.

### **Nanoparticles in sunscreens and cosmetics could cause skin damage**

Scientific studies have shown that nanoparticles of titanium dioxide and zinc oxide commonly used in sunscreens and cosmetics can produce free radicals,<sup>ii</sup> damage DNA<sup>iii</sup> and cause cell toxicity,<sup>iv</sup> especially when exposed to UV light.<sup>v</sup> The concern is that rather than offering us sun protection, nanoparticles used in sunscreens and cosmetics could actually result in serious skin damage.

### ***A world first study shows next generation harm from nanoparticles***

A recent 2009 study by Japanese researchers shows the transfer of nanoparticles of titanium dioxide from pregnant mice to their offspring, with related brain damage, nerve system damage and reduced sperm production in male offspring.<sup>vi</sup> This is concerning because titanium dioxide is one of the most widely used nanoparticles, found in cosmetics, sunscreens, food packaging, paints, wall coatings, dirt repellent coatings for windows, car coatings, etc.

### **No-one knows whether nanoparticles in sunscreen will penetrate intact skin, but it seems likely they could be taken up by damaged or sun burnt skin**

Government agencies worldwide are currently researching whether or not nanoparticles in sunscreens can penetrate intact healthy skin. However scientific studies have shown that nanoparticles not used in sunscreen can penetrate skin,<sup>vii</sup> especially if skin is flexed<sup>viii</sup> (as during exercise), or exposed to penetration enhancers<sup>ix</sup> (as are found in many cosmetics). Damaged skin can take up particles 70 times bigger than a nanoparticle,<sup>x</sup> suggesting that skin penetration by nanoparticles is likely in people with eczema or acne. Recent research has shown that skin penetration is also more likely in sunburnt skin.<sup>xi</sup> We don't know if nanoparticles would more likely penetrate thinner skin – e.g. in elderly people or babies.

### **“Fullerenes” used in cosmetics pose particularly concerning toxic risks and we do know that they penetrate intact skin**

Fullerenes – hollow soccer ball shaped particles made of carbon atoms - are extremely small (0.7nm diameter) so they have very high bioavailability (access to cells etc). Early studies have shown that fullerenes can penetrate intact skin, especially if skin is flexed<sup>xii</sup> or exposed to penetration enhancers.<sup>xiii</sup> Preliminary research suggests that in some forms of fullerenes pose very serious toxicity risks. Carbon fullerenes (buckyballs) have been found to cause brain damage in fish,<sup>xiv</sup> kill water fleas and have bactericidal properties<sup>xv</sup>. Even low levels of exposure to water soluble fullerenes have been shown to be toxic to human liver carcinoma cells and dermal fibroblasts in test tube studies.<sup>xvi</sup> Fullerene-based amino acid nanoparticles have been found to decrease the viability of human epidermal cells and to initiate a pro-inflammatory response.<sup>xvii</sup> Perhaps of most concern, even forms of ‘pristine’ fullerenes that are non-toxic in dark conditions can be highly toxic when exposed to UV light.<sup>xviii</sup>

Cosmetics in which fullerenes are found that are sold in the United States include:

Circuit Addict: <http://www.circuitskin.com/inc/pdetail?v=1&pid=51>

Circuit White Out: <http://www.circuitskin.com/inc/pdetail?v=1&pid=46>

Circuit O.M.G. Serum: <http://www.circuitskin.com/inc/pdetail?v=1&pid=2270>

However with no regulation or labeling in place – it’s impossible to determine the full extent of fullerenes use in cosmetics products.

### **Senior scientists have called for nanoparticles to face new safety testing – what are governments doing?**

The United Kingdom’s Royal Society, the world’s oldest scientific institution, has recommended that given the evidence of serious nanotoxicity risks, nanoparticles should be treated as new chemicals<sup>xix</sup> and subject to new safety assessments before being allowed in consumer products.<sup>xx</sup> Yet, none of the nano-cosmetics on the market are subject to approval by the US Food and Drug Administration (FDA) before being put on the shelves.<sup>xxi</sup> While agencies in the U.S. have hesitantly acknowledged the dangers posed by manufactured nanoparticles, the European Union has taken action to protect the public. New rules passed in Europe on March 2009 will require nanoparticles in cosmetics to be labeled on the ingredients list; the regulation will also require increased safety testing for cosmetics containing nanoparticles.<sup>xxii</sup>

[For more information on nanotechnologies and cosmetics, please refer to two reports by Friends of the Earth U.S. and Australia: “Nanomaterials, Sunscreens and Cosmetics: Small ingredients, Big Risks” and “Nanotechnology and Sunscreens: A Consumer Guide for Avoiding Nano-sunscreens.” These reports can be downloaded at: [www.foe.org](http://www.foe.org)]

## References

---

i <http://www.tga.gov.au/npmeds/sunscreen-zotd.htm>

ii Nel A, Xia T, Li N. 2006. "Toxic potential of materials at the nanolevel". *Science* Vol 311:622-627

iii Donaldson K, Beswick P, Gilmour P. 1996. Free radical activity associated with the surface of particles: a unifying factor in determining biological activity? *Toxicol Lett* 88:293-298.

iv Brunner T, Piusmanser P, Spohn P, Grass R, Limbach L, Bruinink A, Stark W. 2006. In Vitro Cytotoxicity of Oxide Nanoparticles: Comparison to Asbestos, Silica, and the Effect of Particle Solubility. *Environ Sci Technol* 40:4374-4381; Long T, Saleh N, Tilton R, Lowry G, Veronesi B. 2006. Titanium dioxide (P25) produces reactive oxygen species in immortalized brain microglia (BV2): Implications for nanoparticle neurotoxicity. *Environ Sci Technol* 40(14):4346-4352; Sayes C, Wahi R, Kurian P, Liu Y, West J, Ausman K, Warheit D, Colvin V. 2006. Correlating nanoscale titania structure with toxicity: A cytotoxicity and inflammatory response study with human dermal fibroblasts and human lung epithelial cells. *Toxicol Sci* 92(1):174-185.

v Dunford R, Salinaro A, Cai L, Serpone N, Horikoshi S, Hidaka H, Knowland J. 1997. Chemical oxidation and DNA damage catalysed by inorganic sunscreen ingredients. *FEBS Lett* 418:87-90

vi Takeda K, Suzuki K, Ishihara A, Kubo-Irie M, Fujimoto R, Tabata M, Oshio S, Nihei Y, Ihara T, Sugamata M. 2009. Nanoparticles transferred from pregnant mice to their offspring can damage the genital and cranial nerve systems. *J Health Sci* 55(1):95-102. Available at:

vii Ryman-Rasmussen J, Riviere J, Monteiro-Riviere N. 2006. Penetration of intact skin by quantum dots with diverse physicochemical properties. *Toxicol Sci* 91(1):159-165.

viii Rouse J, Yang J, Ryman-Rasmussen J, Barron A, Monteiro-Riviere N. 2007. Effects of mechanical flexion on the penetration of fullerene amino acid derivatized peptide nanoparticles through skin. *Nano Lett* 7(1):155-160; Tinkle S, Antonini J, Roberts J, Salmen R, DePree K, Adkins E. 2003. Skin as a route of exposure and sensitisation in chronic beryllium disease, *Environ Health Perspect* 111:1202-1208.

ix Monteiro-Riviere N, Yang J, Inman A, Ryman-Rasmussen J, Barron A, Riviere J. 2006. Skin penetration of fullerene substituted amino acids and their interactions with human epidermal keratinocytes. *Toxicol* 168 (#827).

x Oberdörster G, Oberdörster E, Oberdörster J. 2005. Nanotoxicology: an emerging discipline from studies of ultrafine particles. *Environ Health Perspect* 113(7):823-839.

xi Mortensen L, Oberdörster G, Pentland A, DeLouise L. 2008. In Vivo Skin Penetration of Quantum Dot Nanoparticles in the Murine Model: The Effect of UVR. *Nano Lett* 8(9):2779-2787

xii Rouse J, Yang J, Ryman-Rasmussen J, Barron A, Monteiro-Riviere N. 2007. Effects of mechanical flexion on the penetration of fullerene amino acid derivatized peptide nanoparticles through skin. *Nano Lett* 7(1):155-160

xiii Monteiro-Riviere N, Yang J, Inman A, Ryman-Rasmussen J, Barron A, Riviere J. 2006. Skin penetration of fullerene substituted amino acids and their interactions with human epidermal keratinocytes. *Toxicol* 168 (#827).

xiv Oberdörster E. 2004. "Manufactured nanomaterials (fullerenes, C60) induce oxidative stress in the brain of juvenile largemouth bass". *Environmental Health Perspectives* 112:1058-1062

xv Fortner J, Lyon D, Sayes C, Boyd A, Falkner J, Hotze E, Alemany L, Tao Y, Guo W, Ausman K, Colvin V, Hughes J. 2005. "C60 in Water: Nanocrystal Formation and Microbial Response". *Environmental Science and Toxicology* 39(11); 4307-4316

---

xvi Sayes C, Fortner J, Guo W, Lyon D, Boyd A, Ausman K, Tao Y, Sitharaman B, Wilson L, Hughes J, West J, Colvin V. 2004. "The differential cytotoxicity of water-soluble fullerenes". *Nanolett.* 4, 1881-1887

xvii Rouse J, Yang J, Barron A, Monteiro-Riviere N. 2006. "Fullerene-based amino acid nanoparticle interactions with human epidermal keratinocytes". *Toxicology In Vitro*

xviii Kolosnjaj J, Szwarc H, Moussa F. 2007. Toxicity studies of fullerenes and derivatives. *Adv Exp Med Biol.* 620:168-80

xix P85 Recommendation 10, The Royal Society and The Royal Academy of Engineering, UK (2004). Nanoscience and nanotechnologies. Available at <http://www.royalsoc.ac.uk/>

xx P86 Recommendation 12 (i), The Royal Society and The Royal Academy of Engineering, UK (2004). Nanoscience and nanotechnologies. Available at <http://www.royalsoc.ac.uk/>

xxi CFSAN/Office of Cosmetics and Colors. 2003. FDA Authority Over Cosmetics. <http://www.cfsan.fda.gov/~dms/cos-206.html>.

xxii See European Parliament website [http://www.europarl.europa.eu/news/expert/infopress\\_page/066-52333-082-03-13-911-20090323IPR52331-23-03-2009-2009-true/default\\_en.htm](http://www.europarl.europa.eu/news/expert/infopress_page/066-52333-082-03-13-911-20090323IPR52331-23-03-2009-2009-true/default_en.htm)